

PHYTOREMEDIATION RESEARCH ON TNT DEGRADATION

Victor F. Medina

AUTHOR: National Research Council Research Associate, c/o USEPA National Exposure Environmental Research Laboratory, Athens, GA 30605 (706) 355 8226. Fax (706) 355 8440. MEDINA.VIC @ epamail.epa.gov

REFERENCE: *Proceedings of the 1997 Georgia Water Resources Conference*, held March 20-22, 1997 at The University of Georgia, Kathryn J. Hatcher, Editor, Institute of Ecology, The University of Georgia, Athens, Georgia.

INTRODUCTION AND BACKGROUND

Phytoremediation is a remediation technology in which plants are used to treat contaminated soil, groundwater or wastewater. Plants can be used the following ways to assist in remediation;

- By enhanced bacterial and fungal activity in the root zone (Anderson et al., 1993; Cunningham et al., 1995; Schwab et al., 1995).
- By accumulating the contaminant in the stems of the plants which can be harvested (Dushenkov et al. 1995; Nanda Kumar et al., 1995; Moffat, 1995; Kelley and Guerin 1995; Cornish et al., 1995; Wang et al., 1995),
- As an organic "pump" to control groundwater migration (Gatliff, 1994; Licht, 1995),
- By destruction by plant enzymes and other plant proteins (Dec and Bollag, 1994, Terry, 1995, Strand et al., 1995, Carreira and Wolfe, 1997, Medina, et al., 1997).

Plants and microorganisms often have symbiotic relationships, making the root zone, or rhizosphere, a very active area of microbiological activity. Plants can help moderate the environment in the root zone, providing ideal conditions for bacteria and fungi to degrade organic pollutants.

Certain plants hyperaccumulate metals, radionuclides or organic compounds. These plants can be placed in areas of contamination where they remove the contaminant. Afterwards, the plants can be harvested and the contaminant can be disposed, destroyed or recovered.

Many plants require large amounts of water in their daily lives. For example, willow trees may use up to 5,000 gallons per day (Gatliff, 1994). These plants can be used to inexpensively "pump" groundwater in certain, shallow aquifer conditions.

Some plants contain enzymes that can destroy contaminants. Of particular interest are;

- dehalogenase: for destroying chlorinated compounds.
- peroxidase: for destroying phenolic compounds.
- nitroreductase: for destroying explosives and other nitrated compounds.
- nitrilase: for destroying cyanated aromatic compounds.
- phosphatase: for destroying organophosphates.

Phytoremediation is attracting much attention as a low cost treatment alternative. However, phytoremediation has several limitations that must be considered. *In situ* phytoremediation is limited to shallow soils or aquifers. Phytotreatment is likely to be slower than excavation, incineration, adsorption or chemical treatments. Phytoremediation is new and, therefore, regulatory

agencies are not yet familiar with it. There are possibilities of increased contaminant migration with improperly designed phytotreatment. Although there are some effective, full-scale phytoremediation projects underway, phytoremediation is still largely in the experimental stage. Winter may slow or stop phytoremediation. Competition with native plants may eliminate the plant chosen for phytoremediation. Phytoremediation designs must be aware of the use of nuisance plants to avoid their propagation.

Overall, phytoremediation is a promising new technology. In some forms, it is already available and proven. New applications are being tested and developed. Phytoremediation, like bioremediation, is relatively slow. Its main advantage is cost savings.

TREATMENT OF EXPLOSIVES

Many military bases and explosives manufacturing facilities are contaminated with munitions. The most prevalent munitions' contaminant is trinitrotoluene (TNT). Other commonly found contaminants include RDX, HMX, Picric Acid, Trinitrobenzene (TNB), and Tertyl.

Numerous studies have documented that plants can uptake and transform TNT from both soil and water (Palazzo and Leggett, 1986; Folsom et al. 1988; Cataldo et al., 1989; Young, 1995; Mueller et al., 1995; Schnoor et al., 1995; Medina and McCutcheon, 1996; Carreira and Wolfe, 1997, Medina et al. 1997). The rates of transformation compare favorably to those of published microbial studies and there appears to be no lag phase (McCutcheon et al., 1995). Previous work conducted by USEPA/NERL isolated a nitroreductase enzyme in plants (Carreira and Wolfe, 1997). This enzyme is capable of sequentially reducing TNT to triaminotoluene (TAT). TAT is an unstable molecule that autooxidizes in aerobic conditions. However, many plants also contain the laccase enzyme, which catalyzes the breakup of the TAT structure. This mechanism is a promising technology for treating TNT contaminated groundwater.

Plant screening studies were conducted for the treatment of TNT contaminated groundwater (Medina, et al., 1997). Parrot Feather (*Myriophyllum aquaticum*) has been chosen as the plant to concentrate experiments on. Removal rates were found to increase with increasing plant density. Temperature variation studies indicated that as temperature increased, the removal rates of increased until the high temperatures became lethal to the plant.

Some preliminary modeling was conducted for the treatment of TNT and its breakdown products (Medina and McCutcheon, 1996). The models assumed idealized plug-flow, first order kinetics and non-reversible, sequential reactions. Bench-scale, continuous flow through reactor studies were also conducted. High removal percentages were achieved on influent concentrations of 4.7 ppm and 12.1 ppm with a 5-day residence time (Medina et al., 1997).

LITERATURE CITED

- Anderson, T.A., Gutherie, E.A., and Walton, B.T. 1993. Bioremediation in the Rhizosphere. *Environmental Science & Technology*. 27(13):2630-2636.
- Carreira, L.H. and Wolfe, N.L. 1997. Isolation of a Sediment Nitroreductase: Characterization, Antibody Production, and the Use of Antibodies to Determine Possible Sources. *Environmental Chemistry and Toxicology*. Paper in Internal Review.
- Cataldo, D.A., Harvey, S.D., Fellows, R.J., Means, R.M., and McVeety, B.D. 1989. *An Evaluation of the Environmental Fate and Behavior of Munitions Material (TNT, RDX) in Soil and Plant Systems*. Richland, WA: Pacific Northwest Laboratory.
- Cornish, J.E., Goldberg, W.C., Levine, R.S. and Benemann, J.R. 1995. Phytoremediation of Soils Contaminated with Toxic Elements and Radionuclides. In: *Bioremediation of Inorganics*. (Hinchee, R.E., Means, J.L., and Burris, D.R., Eds.). Columbus, OH. Battelle Press.
- Cunningham, S.D., Berti, W.R., and Huang, J.W. 1995. Remediation of Contaminated Soils and Sludges by Green Plants. In: *Bioremediation of Inorganics*. (Hinchee, R.E., Means, J.L., and Burris, D.R., Eds.). Columbus, OH. Battelle Press.
- Dec J. and Bollag, J.M. 1994. Use of Plant Material for the Decontamination of Water Polluted with Phenols. *Biotechnology and Bioengineering*. 44:1132-1139.
- Dushenkov, V., Nanda Kumar, P.B.A., Motto, H., and Raskin, I. 1995. Rhizofiltration: The Use of Plants to Remove Heavy Metals from Aqueous Streams. *Environmental Science and Technology*. 29(5):1239-1245.
- Folsom, B.L., Pennington, J.D., Teeter, C.L., Barton, M.R., and Bright, J.A. *Effects of Soil pH and Treatment Level of Persistence and Plant Uptake of 2,4,6-Trinitrotoluene*. Technical Report EL-88-22. Vicksburg, MS. U.S. Army Engineer Waterways Experiment Station.
- Garliff, E.G. 1994. Vegetative Remediation Process Offers Advantages Over Traditional Pump-and-Treat Technologies. *Remediation*. 4(3):343-352.
- Kelley, R.J., and Guerin, T.F. 1995. Feasibility of Using Hyperaccumulating Plants to Bioremediate Metal-Contaminated Soil. In: *Bioremediation of Inorganics*. (Hinchee, R.E., Means, J.L., and Burris, D.R., Eds.). Columbus, OH. Battelle Press.
- Licht, L. (1995): Perennial Plant Systems Using Poplar Trees for Managing Priority Pollutants at Landfills and Industrial Sites. In: *Emerging Technologies in Hazardous Waste Management VIII. 1995 Extended Abstracts for the Special Symposium*. (Teddard, D.W., Ed.). Atlanta, Georgia. American Chemical Society, Industrial and Engineering Division.
- Medina, V.F., and McCutcheon, S.C. 1996. Phytoremediation: Modeling Removal of TNT and Its Breakdown Products. *Remediation*. 6(4):31-45.
- Medina, V.F., Larson, S.L., Ou, T.Y., McCutcheon, S.C., and Wolfe, N.L. 1997. Engineering Studies of Phytoremediation of TNT-Contaminated Water. Submitted for Review to *Bioremediation Journal*.
- Moffat, A.S. 1995. Plants Proving Worth in Toxic Metal Cleanup. *Science*. 269:302-303.
- Mueller, W.F., Bedell, G.W., Shojaei, S., and Jackson, P.J. (1995): Bioremediation of TNT Wastes by Higher Plants. Kansas State Proceedings.
- Nanda Kumar, P.B.A., Dushenkov, V., Motto, H., and Raskin, I. 1995. Phytoextraction: The use of plants to remove heavy metals from soil. *Environmental Science & Technology*. 29(5):1232-1238.
- Palazzo, A.J. and Leggett, D.C. 1986. "Effect and Disposition of TNT in a Terrestrial Plant." *Journal of Environmental Quality*. 15:49-52.
- Schnoor, J.L., Licht, L.A., McCutcheon, S.C., Wolfe, N.L., and Carreira, L.H. 1995. Phytoremediation of Organic and Nutrient Contaminants. *Environmental Science & Technology*. 29:318A-323A.
- Schwab, A.P., M.K. Banks and M. Arunachalam (1995): Biodegradation of Polycyclic Aromatic Hydrocarbons in Rhizosphere Soil. In: *Bioremediation of Recalcitrant Organics*. (Hinchee, R.E., Anderson, D.B., and Hoeppe, R.E., Eds.). Columbus, OH. Battelle Press.
- Strand, S.E., Newman, L., Ruszaj, M., Wilmoth, J., Shurtleff, B., Brandt, M., Choe, N., Ekuan, G., Duffy, J., Massman, J.W., Heilman, P.E., and Gordon, M.P. 1995. Removal of Trichloroethylene from Aquifers Using Trees. In: *Innovative Technologies for Site Remediation and Hazardous Waste Management, Proceedings of the National Conference of the Environmental Engineering Division of the American Society of Civil Engineers*. Pittsburgh, PA. (Vidic, R.D., and Pohland, F.G., Eds.). New York, NY. ASCE.
- Terry, N., Zayed, A., and Pilon-Smits, E. 1995. Bioremediation of Selenium by Plant Volatilization. In: *Emerging Technologies in Hazardous Waste Management VIII. 1995 Extended Abstracts for the Special Symposium*. (Teddard, D.W., Ed.). Atlanta, Georgia. American Chemical Society, Industrial and Engineering Division.
- Wang, T.C., Weissman, J.C., Ramesh, G., Varadarajan, R., and Benemann, J.R. 1995. Bioremoval of Toxic Elements with Aquatic Plants and Algae. In: *Bioremediation of Inorganics*. (Hinchee, R.E., Means, J.L., and Burris, D.R., Eds.) Columbus, OH. Battelle Press.
- Young, D.G. (1995): *The Biochemical Remediation of TNT Contaminated Soil*. Ph.D. Dissertation. Auburn University.